

Package ‘HSAUR’

October 12, 2022

Title A Handbook of Statistical Analyses Using R (1st Edition)

Date 2022-04-25

Version 1.3-10

Author Brian S. Everitt and Torsten Hothorn

Maintainer Torsten Hothorn <Torsten.Hothorn@R-project.org>

Description Functions, data sets, analyses and examples from the book "A Handbook of Statistical Analyses Using R" (Brian S. Everitt and Torsten Hothorn, Chapman & Hall/CRC, 2006). The first chapter of the book, which is entitled "An Introduction to R", is completely included in this package, for all other chapters, a vignette containing all data analyses is available.

Depends R (>= 2.2.0), tools

Suggests lattice, MASS, scatterplot3d (>= 0.3-23), ape (>= 1.6), coin (>= 1.1-3), flexmix (>= 1.1-0), gee (>= 4.13-10), lme4 (>= 0.98-1), mclust (>= 3.0-0), party (>= 0.2-8), randomForest (>= 4.5-12), rmeta (>= 2.12), vcd (>= 0.9-3), survival, KernSmooth, rpart, mvtnorm, Matrix, boot, TH.data

LazyData yes

License GPL

NeedsCompilation no

Repository CRAN

Date/Publication 2022-04-26 04:52:04 UTC

R topics documented:

agefat	2
aspirin	3
BCG	4
birthdeathrates	5
bladdercancer	6
BtheB	7
clouds	8

CYGOB1	9
epilepsy	10
Forbes2000	11
foster	12
gardenflowers	13
GHQ	13
heptathlon	14
HSAURtable	16
Lanza	17
mastectomy	18
meteo	19
orallesions	20
phosphate	20
pistonrings	21
planets	22
plasma	23
polyps	24
polyps3	25
pottery	26
rearrests	27
respiratory	27
roomwidth	28
schizophrenia	29
schizophrenia2	30
schooldays	31
skulls	32
smoking	33
students	34
suicides	35
toothpaste	35
voting	36
water	37
watervoles	38
waves	39
weightgain	40
womensrole	41
Index	42

 agefat

Total Body Composition Data

Description

Age and body fat percentage of 25 normal adults.

Usage

```
data("agefat")
```

Format

A data frame with 25 observations on the following 3 variables.

age the age of the subject.

fat the body fat percentage.

sex a factor with levels female and male.

Details

The data come from a study investigating a new methods of measuring body composition (see Mazess et al, 1984), and give the body fat percentage (percent fat), age and sex for 25 normal adults aged between 23 and 61 years. The questions of interest are how are age and percent fat related, and is there any evidence that the relationship is different for males and females.

Source

R. B. Mazess, W. W. Peppler and M. Gibbons (1984), Total body composition by dual-photon (153Gd) absorptiometry. *American Journal of Clinical Nutrition*, **40**, 834–839.

Examples

```
data("agefat", package = "HSAUR")
plot(fat ~ age, data = agefat)
```

aspirin

Aspirin Data

Description

Efficacy of Aspirin in preventing death after a myocardial infarct.

Usage

```
data("aspirin")
```

Format

A data frame with 7 observations on the following 4 variables.

dp number of deaths after placebo.

tp total number subjects treated with placebo.

da number of deaths after Aspirin.

ta total number of subjects treated with Aspirin.

Details

The data were collected for a meta-analysis of the effectiveness of Aspirin (versus placebo) in preventing death after a myocardial infarction.

Source

J. L. Fleiss (1993), The statistical basis of meta-analysis. *Statistical Methods in Medical Research* **2**, 121–145.

Examples

```
data("aspirin", package = "HSAUR")
aspirin
```

 BCG

BCG Vaccine Data

Description

A meta-analysis on the efficacy of BCG vaccination against tuberculosis (TB).

Usage

```
data("BCG")
```

Format

A data frame with 13 observations on the following 7 variables.

Study an identifier of the study.

BCGTB the number of subjects suffering from TB after a BCG vaccination.

BCGVacc the number of subjects with BCG vaccination.

NoVaccTB the number of subjects suffering from TB without BCG vaccination.

NoVacc the total number of subjects without BCG vaccination.

Latitude geographic position of the place the study was undertaken.

Year the year the study was undertaken.

Details

Bacille Calmette Guerin (BCG) is the most widely used vaccination in the world. Developed in the 1930s and made of a live, weakened strain of *Mycobacterium bovis*, the BCG is the only vaccination available against tuberculosis today. Colditz et al. (1994) report data from 13 clinical trials of BCG vaccine each investigating its efficacy in the treatment of tuberculosis. The number of subjects suffering from TB with or without BCG vaccination are given here. In addition, the data contains the values of two other variables for each study, namely, the geographic latitude of the place where the study was undertaken and the year of publication. These two variables will be used to investigate and perhaps explain any heterogeneity among the studies.

Source

G. A. Colditz, T. F. Brewer, C. S. Berkey, M. E. Wilson, E. Burdick, H. V. Fineberg and F. Mosteller (1994), Efficacy of BCG vaccine in the prevention of tuberculosis. Meta-analysis of the published literature. *Journal of the American Medical Association*, **271**(2), 698–702.

Examples

```
data("BCG", package = "HSAUR")
boxplot(BCG$BCGTB/BCG$BCGVacc, BCG$NoVaccTB/BCG$NoVacc,
        names = c("BCG Vaccination", "No Vaccination"),
        ylab = "Percent BCG cases")
```

birthdeathrates

Birth and Death Rates Data

Description

Birth and death rates for 69 countries.

Usage

```
data("birthdeathrates")
```

Format

A data frame with 69 observations on the following 2 variables.

birth birth rate.

death death rate.

Source

J. A. Hartigan (1975), *Clustering Algorithms*. John Wiley & Sons, New York.

Examples

```
data("birthdeathrates", package = "HSAUR")
plot(birthdeathrates)
```

bladdercancer

Bladder Cancer Data

Description

Data arise from 31 male patients who have been treated for superficial bladder cancer, and give the number of recurrent tumours during a particular time after the removal of the primary tumour, along with the size of the original tumour.

Usage

```
data("bladdercancer")
```

Format

A data frame with 31 observations on the following 3 variables.

time the duration.

tumorsize a factor with levels ≤ 3 cm and > 3 cm.

number number of recurrent tumours.

Details

The aim is to estimate the effect of size of tumour on the number of recurrent tumours.

Source

G. U. H. Seeber (1998), Poisson Regression. In: *Encyclopedia of Biostatistics* (P. Armitage and T. Colton, eds), John Wiley & Sons, Chichester.

Examples

```
data("bladdercancer", package = "HSAUR")  
mosaicplot(xtabs(~ number + tumorsize, data = bladdercancer))
```

BtheB

Beat the Blues Data

Description

Data from a clinical trial of an interactive multimedia program called 'Beat the Blues'.

Usage

```
data("BtheB")
```

Format

A data frame with 100 observations of 100 patients on the following 8 variables.

drug did the patient take anti-depressant drugs (No or Yes).

length the length of the current episode of depression, a factor with levels <6m (less than six months) and >6m (more than six months).

treatment treatment group, a factor with levels TAU (treatment as usual) and BtheB (Beat the Blues)

bdi.pre Beck Depression Inventory II before treatment.

bdi.2m Beck Depression Inventory II after two months.

bdi.4m Beck Depression Inventory II after four months.

bdi.6m Beck Depression Inventory II after six months.

bdi.8m Beck Depression Inventory II after eight months.

Details

Longitudinal data from a clinical trial of an interactive, multimedia program known as "Beat the Blues" designed to deliver cognitive behavioural therapy to depressed patients via a computer terminal. Patients with depression recruited in primary care were randomised to either the Beating the Blues program, or to "Treatment as Usual (TAU)".

Note that the data are stored in the wide form, i.e., repeated measurements are represented by additional columns in the data frame.

Source

J. Proudfoot, D. Goldberg and A. Mann (2003). Computerised, interactive, multimedia CBT reduced anxiety and depression in general practice: A RCT. *Psychological Medicine*, **33**, 217–227.

Examples

```
data("BtheB", package = "HSAUR")
layout(matrix(1:2, nrow = 1))
ylim <- range(BtheB[,grep("bdi", names(BtheB))], na.rm = TRUE)
boxplot(subset(BtheB, treatment == "TAU")[,grep("bdi", names(BtheB))],
        main = "Treated as usual", ylab = "BDI",
        xlab = "Time (in months)", names = c(0, 2, 4, 6, 8), ylim = ylim)
boxplot(subset(BtheB, treatment == "BtheB")[,grep("bdi", names(BtheB))],
        main = "Beat the Blues", ylab = "BDI", xlab = "Time (in months)",
        names = c(0, 2, 4, 6, 8), ylim = ylim)
```

clouds

Cloud Seeding Data

Description

Data from an experiment investigating the use of massive amounts of silver iodide (100 to 1000 grams per cloud) in cloud seeding to increase rainfall.

Usage

```
data("clouds")
```

Format

A data frame with 24 observations on the following 7 variables.

seeding a factor indicating whether seeding action occurred (no or yes).

time number of days after the first day of the experiment.

sne suitability criterion.

cloudcover the percentage cloud cover in the experimental area, measured using radar.

prewetness the total rainfall in the target area one hour before seeding (in cubic metres times $1e+8$).

echomotion a factor showing whether the radar echo was moving or stationary.

rainfall the amount of rain in cubic metres times $1e+8$.

Details

Weather modification, or cloud seeding, is the treatment of individual clouds or storm systems with various inorganic and organic materials in the hope of achieving an increase in rainfall. Introduction of such material into a cloud that contains supercooled water, that is, liquid water colder than zero Celsius, has the aim of inducing freezing, with the consequent ice particles growing at the expense of liquid droplets and becoming heavy enough to fall as rain from clouds that otherwise would produce none.

The data available in `cCloud` were collected in the summer of 1975 from an experiment to investigate the use of massive amounts of silver iodide 100 to 1000 grams per cloud) in cloud seeding to increase rainfall. In the experiment, which was conducted in an area of Florida, 24 days were judged suitable for seeding on the basis that a measured suitability criterion (SNE).

Source

W. L. Woodley, J. Simpson, R. Biondini and J. Berkeley (1977), Rainfall results 1970-75: Florida area cumulus experiment. *Science* **195**, 735–742.

R. D. Cook and S. Weisberg (1980), Characterizations of an empirical influence function for detecting influential cases in regression. *Technometrics* **22**, 495–508.

Examples

```
data("clouds", package = "HSAUR")
layout(matrix(1:2, nrow = 2))
boxplot(rainfall ~ seeding, data = clouds, ylab = "Rainfall")
boxplot(rainfall ~ echomotion, data = clouds, ylab = "Rainfall")
```

CYGOB1

CYG OB1 Star Cluster Data

Description

Energy output and surface temperature for Star Cluster CYG OB1.

Usage

```
data("CYGOB1")
```

Format

A data frame with 47 observations on the following 2 variables.

`logst` log surface temperature of the star.

`logli` log light intensity of the star.

Details

The Hertzsprung-Russell (H-R) diagram forms the basis of the theory of stellar evolution. The diagram is essentially a plot of the energy output of stars plotted against their surface temperature. Data from the H-R diagram of Star Cluster CYG OB1, calibrated according to VanismaGreve1972 are given here.

Source

F. Vanisma and J. P. De Greve (1972), Close binary systems before and after mass transfer. *Astrophysics and Space Science*, **87**, 377–401.

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("CYGOB1", package = "HSAUR")
plot(logst ~ logli, data = CYGOB1)
```

epilepsy

Epilepsy Data

Description

A randomised clinical trial investigating the effect of an anti-epileptic drug.

Usage

```
data("epilepsy")
```

Format

A data frame with 236 observations on the following 6 variables.

`treatment` the treatment group, a factor with levels placebo and Progabide.

`base` the number of seizures before the trial.

`age` the age of the patient.

`seizure.rate` the number of seizures (response variable).

`period` treatment period, an ordered factor with levels 1 to 4.

`subject` the patient ID, a factor with levels 1 to 59.

Details

In this clinical trial, 59 patients suffering from epilepsy were randomized to groups receiving either the anti-epileptic drug Progabide or a placebo in addition to standard chemotherapy. The numbers of seizures suffered in each of four, two-week periods were recorded for each patient along with a baseline seizure count for the 8 weeks prior to being randomized to treatment and age. The main question of interest is whether taking progabide reduced the number of epileptic seizures compared with placebo.

Source

P. F. Thall and S. C. Vail (1990), Some covariance models for longitudinal count data with overdispersion. *Biometrics*, **46**, 657–671.

Examples

```
data("epilepsy", package = "HSAUR")
library(lattice)
dotplot(I(seizure.rate / base) ~ period | subject, data = epilepsy,
        subset = treatment == "Progabide")
dotplot(I(seizure.rate / base) ~ period | subject, data = epilepsy,
        subset = treatment == "Progabide")
```

Forbes2000

The Forbes 2000 Ranking of the World's Biggest Companies (Year 2004)

Description

The Forbes 2000 list is a ranking of the world's biggest companies, measured by sales, profits, assets and market value.

Usage

```
data("Forbes2000")
```

Format

A data frame with 2000 observations on the following 8 variables.

rank the ranking of the company.

name the name of the company.

country a factor giving the country the company is situated in.

category a factor describing the products the company produces.

sales the amount of sales of the company in billion USD.

profits the profit of the company in billion USD.

assets the assets of the company in billion USD.

marketvalue the market value of the company in billion USD.

Source

<https://www.forbes.com>, assessed on November 26th, 2004.

Examples

```
data("Forbes2000", package = "HSAUR")
summary(Forbes2000)
### number of countries
length(levels(Forbes2000$country))
### number of industries
length(levels(Forbes2000$category))
```

foster

Foster Feeding Experiment

Description

The data are from a foster feeding experiment with rat mothers and litters of four different genotypes. The measurement is the litter weight after a trial feeding period.

Usage

```
data("foster")
```

Format

A data frame with 61 observations on the following 3 variables.

`litgen` genotype of the litter, a factor with levels A, B, I, and J.

`motgen` genotype of the mother, a factor with levels A, B, I, and J.

`weight` the weight of the litter after a feeding period.

Details

Here the interest lies in uncovering the effect of genotype of mother and litter on litter weight.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("foster", package = "HSAUR")
plot.design(foster)
```

gardenflowers

Garden Flowers

Description

The dissimilarity matrix of 18 species of garden flowers.

Usage

```
data("gardenflowers")
```

Format

An object of class `dist`.

Details

The dissimilarity was computed based on certain characteristics of the flowers.

Source

L. Kaufman and P. J. Rousseeuw (1990), *Finding groups in data: an introduction to cluster analysis*, John Wiley & Sons, New York.

Examples

```
data("gardenflowers", package = "HSAUR")
gardenflowers
```

GHQ

General Health Questionnaire

Description

Data from an psychiatric screening questionnaire

Usage

```
data("GHQ")
```

Format

A data frame with 22 observations on the following 4 variables.

GHQ the General Health Questionnaire score.

sex a factor with levels female and male

cases the number of diseased subjects.

non.cases the number of healthy subjects.

Details

The data arise from a study of a psychiatric screening questionnaire called the GHQ (General Health Questionnaire, see Goldberg, 1972). Here the main question of interest is to see how caseness is related to gender and GHQ score.

Source

D. Goldberg (1972). *The Detection of Psychiatric Illness by Questionnaire*, Oxford University Press, Oxford, UK.

Examples

```
data("GHQ", package = "HSAUR")
male <- subset(GHQ, sex == "male")
female <- subset(GHQ, sex == "female")
layout(matrix(1:2, ncol = 2))
barplot(t(as.matrix(male[,c("cases", "non.cases")])), main = "Male", xlab = "GHC score")
barplot(t(as.matrix(female[,c("cases", "non.cases")])), main = "Female", xlab = "GHC score")
```

heptathlon

Olympic Heptathlon Seoul 1988

Description

Results of the olympic heptathlon competition, Seoul, 1988.

Usage

```
data("heptathlon")
```

Format

A data frame with 25 observations on the following 8 variables.

hurdles results 100m hurdles.

highjump results high jump.

shot results shot.

run200m results 200m race.

longjump results long jump.

javelin results javelin.

run800m results 800m race.

score total score.

Details

The first combined Olympic event for women was the pentathlon, first held in Germany in 1928. Initially this consisted of the shot putt, long jump, 100m, high jump and javelin events held over two days. The pentathlon was first introduced into the Olympic Games in 1964, when it consisted of the 80m hurdles, shot, high jump, long jump and 200m. In 1977 the 200m was replaced by the 800m and from 1981 the IAAF brought in the seven-event heptathlon in place of the pentathlon, with day one containing the events-100m hurdles, shot, high jump, 200m and day two, the long jump, javelin and 800m. A scoring system is used to assign points to the results from each event and the winner is the woman who accumulates the most points over the two days. The event made its first Olympic appearance in 1984.

In the 1988 Olympics held in Seoul, the heptathlon was won by one of the stars of women's athletics in the USA, Jackie Joyner-Kersey. The results for all 25 competitors are given here.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("heptathlon", package = "HSAUR")
plot(heptathlon)
```

HSAURtable

Produce LaTeX Tables

Description

Generate longtable LaTeX environments.

Usage

```
HSAURtable(object, ...)
## S3 method for class 'table'
HSAURtable(object, xname = deparse(substitute(object)), pkg = NULL,
  ...)
## S3 method for class 'data.frame'
HSAURtable(object, xname = deparse(substitute(object)), pkg = NULL,
  nrows = NULL, ...)
## S3 method for class 'tabtab'
toLatex(object, caption = NULL, label = NULL,
  topcaption = TRUE, index = TRUE, ...)
## S3 method for class 'dftab'
toLatex(object, pcol = 1, caption = NULL,
  label = NULL, rownames = FALSE, topcaption = TRUE, index = TRUE,
  ...)
```

Arguments

object	an object of table or data.frame.
xname	the name of the object.
pkg	the package object comes from, optionally.
nrows	the number of rows actually printed for a data.frame.
caption	the (optional) caption of the table without label.
label	the (optional) label to be defined for this table.
pcol	the number of parallel columns.
rownames	logical, should the rownames be printed in the first row without column name?
topcaption	logical, should the captions be placed on top (default) of the table?
index	logical, should an index entry be generated?
...	additional arguments, currently ignored.

Details

Based on the data in object, an object from which a Latex table (in a longtable environment) may be constructed (via [toLatex](#)) is generated.

Value

An object of class `tabtab` or `dftab` for which `toLatex` methods are available.
`toLatex` produces objects of class `Latex`, a character vector, essentially.

Examples

```
data("rearrests", package = "HSAUR")
toLatex(HSAURtable(rearrests),
        caption = "Rearrests of juvenile felons.",
        label = "rearrests_tab")
```

Lanza

Prevention of Gastrointestinal Damages

Description

Data from four randomised clinical trials on the prevention of gastrointestinal damages by Misoprostol reported by Lanza et al. (1987, 1988a,b, 1989).

Usage

```
data("Lanza")
```

Format

A data frame with 198 observations on the following 3 variables.

`study` a factor with levels I, II, III, and IV describing the study number.

`treatment` a factor with levels Misoprostol Placebo

`classification` an ordered factor with levels 1 < 2 < 3 < 4 < 5 describing an ordered response variable.

Details

The response variable is defined by the number of haemorrhages or erosions.

Source

F. L. Lanza (1987), A double-blind study of prophylactic effect of misoprostol on lesions of gastric and duodenal mucosa induced by oral administration of tolmetin in healthy subjects. *British Journal of Clinical Practice*, May suppl, 91–101.

F. L. Lanza, R. L. Aspinall, E. A. Swabb, R. E. Davis, M. F. Rack, A. Rubin (1988a), Double-blind, placebo-controlled endoscopic comparison of the mucosal protective effects of misoprostol versus cimetidine on tolmetin-induced mucosal injury to the stomach and duodenum. *Gastroenterology*, **95**(2), 289–294.

F. L. Lanza, K. Peace, L. Gustitus, M. F. Rack, B. Dickson (1988b), A blinded endoscopic comparative study of misoprostol versus sucralfate and placebo in the prevention of aspirin-induced gastric and duodenal ulceration. *American Journal of Gastroenterology*, **83**(2), 143–146.

F. L. Lanza, D. Fakouhi, A. Rubin, R. E. Davis, M. F. Rack, C. Nissen, S. Geis (1989), A double-blind placebo-controlled comparison of the efficacy and safety of 50, 100, and 200 micrograms of misoprostol QID in the prevention of ibuprofen-induced gastric and duodenal mucosal lesions and symptoms. *American Journal of Gastroenterology*, **84**(6), 633–636.

Examples

```
data("Lanza", package = "HSAUR")
layout(matrix(1:4, nrow = 2))
pl <- tapply(1:nrow(Lanza), Lanza$study, function(indx)
  mosaicplot(table(Lanza[indx,"treatment"],
    Lanza[indx,"classification"]),
    main = "", shade = TRUE))
```

mastectomy

Survival Times after Mastectomy of Breast Cancer Patients

Description

Survival times in months after mastectomy of women with breast cancer. The cancers are classified as having metastized or not based on a histochemical marker.

Usage

```
data("mastectomy")
```

Format

A data frame with 42 observations on the following 3 variables.

time survival times in months.

event a logical indicating if the event was observed (TRUE) or if the survival time was censored (FALSE).

metastized a factor at levels yes and no.

Source

B. S. Everitt and S. Rabe-Hesketh (2001), *Analysing Medical Data using S-PLUS*, Springer, New York, USA.

Examples

```
data("mastectomy", package = "HSAUR")
table(mastectomy$metastized)
```

meteo

Meteorological Measurements for 11 Years

Description

Several meteorological measurements for a period between 1920 and 1931.

Usage

```
data("meteo")
```

Format

A data frame with 11 observations on the following 6 variables.

year the years.

rainNovDec rainfall in November and December (mm).

temp average July temperature.

rainJuly rainfall in July (mm).

radiation radiation in July (millilitres of alcohol).

yield average harvest yield (quintals per hectare).

Details

Carry out a principal components analysis of both the covariance matrix and the correlation matrix of the data and compare the results. Which set of components leads to the most meaningful interpretation?

Source

B. S. Everitt and G. Dunn (2001), *Applied Multivariate Data Analysis*, 2nd edition, Arnold, London.

Examples

```
data("meteo", package = "HSAUR")  
meteo
```

orallesions

Oral Lesions in Rural India

Description

The distribution of the oral lesion site found in house-to-house surveys in three geographic regions of rural India.

Usage

```
data("orallesions")
```

Format

A two-way classification, see [table](#).

Source

Cyrus R. Mehta and Nitin R. Patel (2003), *StatXact-6: Statistical Software for Exact Nonparametric Inference*, Cytel Software Cooperation, Cambridge, USA.

Examples

```
data("orallesions", package = "HSAUR")  
mosaicplot(orallesions)
```

phosphate

Phosphate Level Data

Description

Plasma inorganic phosphate levels from 33 subjects.

Usage

```
data("phosphate")
```

Format

A data frame with 33 observations on the following 9 variables.

group a factor with levels control and obese.

t0 baseline phosphate level,

t0.5 phosphate level after 1/2 an hour.

t1 phosphate level after one an hour.

t1.5 phosphate level after 1 1/2 hours.

t2 phosphate level after two hours.

t3 phosphate level after three hours.

t4 phosphate level after four hours.

t5 phosphate level after five hours.

Source

C. S. Davis (2002), *Statistical Methods for the Analysis of Repeated Measurements*, Springer, New York.

Examples

```
data("phosphate", package = "HSAUR")
plot(t0 ~ group, data = phosphate)
```

pistonrings

Piston Rings Failures

Description

Number of failures of piston rings in three legs of four steam-driven compressors.

Usage

```
data("pistonrings")
```

Format

A two-way classification, see [table](#).

Details

The data are given in form of a [table](#). The table gives the number of piston-ring failures in each of three legs of four steam-driven compressors located in the same building. The compressors have identical design and are oriented in the same way. The question of interest is whether the two classification variables (compressor and leg) are independent.

Source

S. J. Haberman (1973), The analysis of residuals in cross-classified tables. *Biometrics* **29**, 205–220.

Examples

```
data("pistonrings", package = "HSAUR")
mosaicplot(pistonrings)
```

planets

Exoplanets Data

Description

Data on planets outside the Solar System.

Usage

```
data("planets")
```

Format

A data frame with 101 observations from 101 exoplanets on the following 3 variables.

mass Jupiter mass of the planet.

period period in earth days.

eccen the radial eccentricity of the planet.

Details

From the properties of the exoplanets found up to now it appears that the theory of planetary development constructed for the planets of the Solar System may need to be reformulated. The exoplanets are not at all like the nine local planets that we know so well. A first step in the process of understanding the exoplanets might be to try to classify them with respect to their known properties.

Source

M. Mayor and P. Frei (2003). *New Worlds in the Cosmos: The Discovery of Exoplanets*. Cambridge University Press, Cambridge, UK.

Examples

```
data("planets", package = "HSAUR")
require("scatterplot3d")
scatterplot3d(log(planets$mass), log(planets$period), log(planets$eccen),
              type = "h", highlight.3d = TRUE, angle = 55,
              scale.y = 0.7, pch = 16)
```

plasma

Blood Screening Data

Description

The erythrocyte sedimentation rate and measurements of two plasma proteins (fibrinogen and globulin).

Usage

```
data("plasma")
```

Format

A data frame with 32 observations on the following 3 variables.

`fibrinogen` the fibrinogen level in the blood.

`globulin` the globulin level in the blood.

`ESR` the erythrocyte sedimentation rate, either less or greater 20 mm / hour.

Details

The erythrocyte sedimentation rate (ESR) is the rate at which red blood cells (erythrocytes) settle out of suspension in blood plasma, when measured under standard conditions. If the ESR increases when the level of certain proteins in the blood plasma rise in association with conditions such as rheumatic diseases, chronic infections and malignant diseases, its determination might be useful in screening blood samples taken from people suspected to be suffering from one of the conditions mentioned. The absolute value of the ESR is not of great importance rather it is whether it is less than 20mm/hr since lower values indicate a healthy individual.

The question of interest is whether there is any association between the probability of an ESR reading greater than 20mm/hr and the levels of the two plasma proteins. If there is not then the determination of ESR would not be useful for diagnostic purposes.

Source

D. Collett and A. A. Jemain (1985), Residuals, outliers and influential observations in regression analysis. *Sains Malaysiana*, **4**, 493–511.

Examples

```
data("plasma", package = "HSAUR")
layout(matrix(1:2, ncol = 2))
boxplot(fibrinogen ~ ESR, data = plasma, varwidth = TRUE)
boxplot(globulin ~ ESR, data = plasma, varwidth = TRUE)
```

polyps

Familial Adenomatous Polyposis

Description

Data from a placebo-controlled trial of a non-steroidal anti-inflammatory drug in the treatment of familial adenomatous polyposis (FAP).

Usage

```
data("polyps")
```

Format

A data frame with 20 observations on the following 3 variables.

number number of colonic polyps at 12 months.

treat treatment arms of the trial, a factor with levels placebo and drug.

age the age of the patient.

Details

Giardiello et al. (1993) and Piantadosi (1997) describe the results of a placebo-controlled trial of a non-steroidal anti-inflammatory drug in the treatment of familial adenomatous polyposis (FAP). The trial was halted after a planned interim analysis had suggested compelling evidence in favour of the treatment. Here we are interested in assessing whether the number of colonic polyps at 12 months is related to treatment and age of patient.

Source

F. M. Giardiello, S. R. Hamilton, A. J. Krush, S. Piantadosi, L. M. Hyland, P. Celano, S. V. Booker, C. R. Robinson and G. J. A. Offerhaus (1993), Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *New England Journal of Medicine*, **328**(18), 1313–1316.

S. Piantadosi (1997), *Clinical Trials: A Methodologic Perspective*. John Wiley & Sons, New York.

Examples

```
data("polyps", package = "HSAUR")
plot(number ~ age, data = polyps, pch = as.numeric(polyps$treat))
legend(40, 40, legend = levels(polyps$treat), pch = 1:2, bty = "n")
```

 polyps3

Familial Adenomatous Polyposis

Description

Data from a placebo-controlled trial of a non-steroidal anti-inflammatory drug in the treatment of familial adenomatous polyposis (FAP).

Usage

```
data("polyps3")
```

Format

A data frame with 22 observations on the following 5 variables.

`sex` a factor with levels female and male.

`treatment` a factor with levels placebo and active.

`baseline` the baseline number of polyps.

`age` the age of the patient.

`number3m` the number of polyps after three month.

Details

The data arise from the same study as the `polyps` data. Here, the number of polyps after three months are given.

Source

F. M. Giardiello, S. R. Hamilton, A. J. Krush, S. Piantadosi, L. M. Hylind, P. Celano, S. V. Booker, C. R. Robinson and G. J. A. Offerhaus (1993), Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *New England Journal of Medicine*, **328**(18), 1313–1316.

S. Piantadosi (1997), *Clinical Trials: A Methodologic Perspective*. John Wiley & Sons, New York.

Examples

```
data("polyps3", package = "HSAUR")
plot(number3m ~ age, data = polyps3, pch = as.numeric(polyps3$treatment))
legend("topright", legend = levels(polyps3$treatment), pch = 1:2, bty = "n")
```

pottery

Romano-British Pottery Data

Description

Chemical composition of Romano-British pottery.

Usage

```
data("pottery")
```

Format

A data frame with 45 observations on the following 9 chemicals.

Al₂O₃ aluminium trioxide.

Fe₂O₃ iron trioxide.

MgO magnesium oxide.

CaO calcium oxide.

Na₂O natrium oxide.

K₂O calium oxide.

TiO₂ titanium oxide.

MnO mangan oxide.

BaO barium oxide.

Details

The data gives the chemical composition of specimens of Romano-British pottery, determined by atomic absorption spectrophotometry, for nine oxides.

Source

A. Tubb and N. J. Parker and G. Nickless (1980), The analysis of Romano-British pottery by atomic absorption spectrophotometry. *Archaeometry*, **22**, 153–171.

Examples

```
data("pottery", package = "HSAUR")  
plot(pottery)
```

rearrests

Rearrests of Juvenile Felons

Description

Rearrests of juvenile felons by type of court in which they were tried.

Usage

```
data("rearrests")
```

Format

A two-way classification, see [table](#).

Details

The data (taken from Agresti, 1996) arise from a sample of juveniles convicted of felony in Florida in 1987. Matched pairs were formed using criteria such as age and the number of previous offences. For each pair, one subject was handled in the juvenile court and the other was transferred to the adult court. Whether or not the juvenile was rearrested by the end of 1988 was then noted. Here the question of interest is whether the true proportions rearrested were identical for the adult and juvenile court assignments?

Source

A. Agresti (1996). *An Introduction to Categorical Data Analysis*. Wiley, New York.

Examples

```
data("rearrests", package = "HSAUR")
rearrests
```

respiratory

Respiratory Illness Data

Description

The respiratory status of patients recruited for a randomised clinical multicenter trial.

Usage

```
data("respiratory")
```

Format

A data frame with 555 observations on the following 7 variables.

centre the study center, a factor with levels 1 and 2.

treatment the treatment arm, a factor with levels placebo and treatment.

sex a factor with levels female and male.

age the age of the patient.

status the respiratory status (response variable), a factor with levels poor and good.

month the month, each patient was examined at months 0, 1, 2, 3 and 4.

subject the patient ID, a factor with levels 1 to 111.

Details

In each of two centres, eligible patients were randomly assigned to active treatment or placebo. During the treatment, the respiratory status (categorised poor or good) was determined at each of four, monthly visits. The trial recruited 111 participants (54 in the active group, 57 in the placebo group) and there were no missing data for either the responses or the covariates. The question of interest is to assess whether the treatment is effective and to estimate its effect.

Note that the data are in long form, i.e, repeated measurements are stored as additional rows in the data frame.

Source

C. S. Davis (1991), Semi-parametric and non-parametric methods for the analysis of repeated measurements with applications to clinical trials. *Statistics in Medicine*, **10**, 1959–1980.

Examples

```
data("respiratory", package = "HSAUR")
mosaicplot(xtabs(~ treatment + month + status, data = respiratory))
```

roomwidth

Students Estimates of Lecture Room Width

Description

Lecture room width estimated by students in two different units.

Usage

```
data("roomwidth")
```

Format

A data frame with 113 observations on the following 2 variables.

unit a factor with levels feet and metres.

width the estimated width of the lecture room.

Details

Shortly after metric units of length were officially introduced in Australia, each of a group of 44 students was asked to guess, to the nearest metre, the width of the lecture hall in which they were sitting. Another group of 69 students in the same room was asked to guess the width in feet, to the nearest foot. The data were collected by Professor T. Lewis and are taken from Hand et al (1994). The main question is whether estimation in feet and in metres gives different results.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("roomwidth", package = "HSAUR")
convert <- ifelse(roomwidth$unit == "feet", 1, 3.28)
boxplot(I(width * convert) ~ unit, data = roomwidth)
```

schizophrenia

Age of Onset of Schizophrenia Data

Description

Data on sex differences in the age of onset of schizophrenia.

Usage

```
data("schizophrenia")
```

Format

A data frame with 251 observations on the following 2 variables.

age age at the time of diagnosis.

gender a factor with levels female and male

Details

A sex difference in the age of onset of schizophrenia was noted by Kraepelin (1919). Subsequently epidemiological studies of the disorder have consistently shown an earlier onset in men than in women. One model that has been suggested to explain this observed difference is known as the subtype model which postulates two types of schizophrenia, one characterised by early onset, typical symptoms and poor premorbid competence, and the other by late onset, atypical symptoms, and good premorbid competence. The early onset type is assumed to be largely a disorder of men and the late onset largely a disorder of women.

Source

E. Kraepelin (1919), *Dementia Praecox and Paraphrenia*. Livingstone, Edinburgh.

Examples

```
data("schizophrenia", package = "HSAUR")
boxplot(age ~ gender, data = schizophrenia)
```

schizophrenia2

Schizophrenia Data

Description

Though disorder and early onset of schizophrenia.

Usage

```
data("schizophrenia2")
```

Format

A data frame with 220 observations on the following 4 variables.

subject the patient ID, a factor with levels 1 to 44.

onset the time of onset of the disease, a factor with levels < 20 yrs and > 20 yrs.

disorder whether thought disorder was absent or present, the response variable.

month month after hospitalisation.

Details

The data were collected in a follow-up study of women patients with schizophrenia. The binary response recorded at 0, 2, 6, 8 and 10 months after hospitalisation was thought disorder (absent or present). The single covariate is the factor indicating whether a patient had suffered early or late onset of her condition (age of onset less than 20 years or age of onset 20 years or above). The question of interest is whether the course of the illness differs between patients with early and late onset?

Source

Davis (2002), *Statistical Methods for the Analysis of Repeated Measurements*, Springer, New York.

Examples

```
data("schizophrenia2", package = "HSAUR")
mosaicplot(xtabs(~ onset + month + disorder, data = schizophrenia2))
```

schooldays

Days not Spent at School

Description

Data from a sociological study, the number of days absent from school is the response variable.

Usage

```
data("schooldays")
```

Format

A data frame with 154 observations on the following 5 variables.

race race of the child, a factor with levels aboriginal and non-aboriginal.

sex the sex of the child, a factor with levels female and male.

school the school type, a factor with levels F0 (primary), F1 (first), F2 (second) and F3 (third form).

learner how good is the child in learning things, a factor with levels average and slow.

absent number of days absent from school.

Details

The data arise from a sociological study of Australian Aboriginal and white children reported by Quine (1975).

In this study, children of both sexes from four age groups (final grade in primary schools and first, second and third form in secondary school) and from two cultural groups were used. The children in age group were classified as slow or average learners. The response variable was the number of days absent from school during the school year. (Children who had suffered a serious illness during the years were excluded.)

Source

S. Quine (1975), *Achievement Orientation of Aboriginal and White Adolescents*. Doctoral Dissertation, Australian National University, Canberra.

Examples

```
data("schooldays", package = "HSAUR")
plot.design(schooldays)
```

 skulls

*Egyptian Skulls***Description**

Measurements made on Egyptian skulls from five epochs.

Usage

```
data("skulls")
```

Format

A data frame with 150 observations on the following 5 variables.

epoch the epoch the skull as assigned to, a factor with levels c4000BC c3300BC, c1850BC, c200BC, and cAD150, where the years are only given approximately, of course.

mb maximum breaths of the skull.

bh basibregmatic heights of the skull.

b1 basialveolar length of the skull.

nh nasal heights of the skull.

Details

The question is whether the measurements change over time. Non-constant measurements of the skulls over time would indicate interbreeding with immigrant populations.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("skulls", package = "HSAUR")
means <- tapply(1:nrow(skulls), skulls$epoch, function(i)
  apply(skulls[i,colnames(skulls)[-1]], 2, mean))
means <- matrix(unlist(means), nrow = length(means), byrow = TRUE)
colnames(means) <- colnames(skulls)[-1]
rownames(means) <- levels(skulls$epoch)
```



```
pairs(means,
      panel = function(x, y) {
        text(x, y, levels(skulls$epoch))
      })
```

smoking

Nicotine Gum and Smoking Cessation

Description

Data from a meta-analysis on nicotine gum and smoking cessation

Usage

```
data("smoking")
```

Format

A data frame with 26 observations (studies) on the following 4 variables.

qt the number of treated subjects who stopped smoking.

tt the total number of treated subjects.

qc the number of subjects who stopped smoking without being treated.

tc the total number of subject not being treated.

Details

Cigarette smoking is the leading cause of preventable death in the United States and kills more Americans than AIDS, alcohol, illegal drug use, car accidents, fires, murders and suicides combined. It has been estimated that 430,000 Americans die from smoking every year. Fighting tobacco use is, consequently, one of the major public health goals of our time and there are now many programs available designed to help smokers quit. One of the major aids used in these programs is nicotine chewing gum, which acts as a substitute oral activity and provides a source of nicotine that reduces the withdrawal symptoms experienced when smoking is stopped. But separate randomized clinical trials of nicotine gum have been largely inconclusive, leading Silagy (2003) to consider combining the results studies found from an extensive literature search. The results of these trials in terms of numbers of people in the treatment arm and the control arm who stopped smoking for at least 6 months after treatment are given here.

Source

C. Silagy (2003), Nicotine replacement therapy for smoking cessation (Cochrane Review). *The Cochrane Library*, 4, John Wiley & Sons, Chichester.

Examples

```
data("smoking", package = "HSAUR")
boxplot(smoking$qt/smoking$tt,
        smoking$qc/smoking$tc,
        names = c("Treated", "Control"), ylab = "Percent Quitters")
```

 students

Student Risk Taking

Description

Students were administered two parallel forms of a test after a random assignment to three different treatments.

Usage

```
data("students")
```

Format

A data frame with 35 observations on the following 3 variables.

treatment a factor with levels AA, C, and NC.

low the result of the first test.

high the result of the second test.

Details

The data arise from a large study of risk taking (Timm, 2002). Students were randomly assigned to three different treatments labelled AA, C and NC. Students were administered two parallel forms of a test called low and high. The aim is to carry out a test of the equality of the bivariate means of each treatment population.

Source

N. H. Timm (2002), *Applied Multivariate Analysis*. Springer, New York.

Examples

```
data("students", package = "HSAUR")
layout(matrix(1:2, ncol = 2))
boxplot(low ~ treatment, data = students, ylab = "low")
boxplot(high ~ treatment, data = students, ylab = "high")
```

suicides

Crowd Baiting Behaviour and Suicides

Description

Data from a study carried out to investigate the causes of jeering or baiting behaviour by a crowd when a person is threatening to commit suicide by jumping from a high building.

Usage

```
data("suicides")
```

Format

A two-way classification, see [table](#).

Source

L. Mann (1981), The baiting crowd in episodes of threatened suicide. *Journal of Personality and Social Psychology*, **41**, 703–709.

Examples

```
data("suicides", package = "HSAUR")  
mosaicplot(suicides)
```

toothpaste

Toothpaste Data

Description

Meta-analysis of studies comparing two different toothpastes.

Usage

```
data("toothpaste")
```

Format

A data frame with 9 observations on the following 7 variables.

Study the identifier of the study.

nA number of subjects using toothpaste A.

meanA mean DMFS index of subjects using toothpaste A.

sdA standard deviation of DMFS index of subjects using toothpaste A.

nB number of subjects using toothpaste B.

meanB mean DMFS index of subjects using toothpaste B.

sdB standard deviation of DMFS index of subjects using toothpaste B.

Details

The data are the results of nine randomised trials comparing two different toothpastes for the prevention of caries development. The outcomes in each trial was the change, from baseline, in the decayed, missing (due to caries) and filled surface dental index (DMFS).

Source

B. S. Everitt and A. Pickles (2000), *Statistical Aspects of the Design and Analysis of Clinical Trials*, Imperial College Press, London.

Examples

```
data("toothpaste", package = "HSAUR")
toothpaste
```

voting

House of Representatives Voting Data

Description

Voting results for 15 congressmen from New Jersey.

Usage

```
data("voting")
```

Format

A 15 times 15 matrix.

Details

Romesburg (1984) gives a set of data that shows the number of times 15 congressmen from New Jersey voted differently in the House of Representatives on 19 environmental bills. Abstentions are not recorded.

Source

H. C. Romesburg (1984), *Cluster Analysis for Researchers*. Lifetime Learning Publications, Belmont, Canada.

Examples

```
data("voting", package = "HSAUR")
require("MASS")
voting_mds <- isoMDS(voting)
plot(voting_mds$points[,1], voting_mds$points[,2],
     type = "n", xlab = "Coordinate 1", ylab = "Coordinate 2",
     xlim = range(voting_mds$points[,1])*1.2)
text(voting_mds$points[,1], voting_mds$points[,2],
     labels = colnames(voting))
voting_sh <- Shepard(voting[lower.tri(voting)], voting_mds$points)
```

water

Mortality and Water Hardness

Description

The mortality and drinking water hardness for 61 cities in England and Wales.

Usage

```
data("water")
```

Format

A data frame with 61 observations on the following 4 variables.

location a factor with levels North and South indicating whether the town is as north as Derby.

town the name of the town.

mortality averaged annual mortality per 100.000 male inhabitants.

hardness calcium concentration (in parts per million).

Details

The data were collected in an investigation of environmental causes of disease. They show the annual mortality per 100,000 for males, averaged over the years 1958-1964, and the calcium concentration (in parts per million) in the drinking water for 61 large towns in England and Wales. The higher the calcium concentration, the harder the water. Towns at least as far north as Derby are identified in the table. Here there are several questions that might be of interest including, are mortality and water hardness related, and do either or both variables differ between northern and southern towns?

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("water", package = "HSAUR")
plot(mortality ~ hardness, data = water,
     col = as.numeric(water$location))
```

 watervoles

Water Voles Data

Description

Percentage incidence of the 13 characteristics of water voles in 14 areas.

Usage

```
data("watervoles")
```

Format

A dissimilarity matrix for the following 14 variables, i.e, areas: Surrey, Shropshire, Yorkshire, Perthshire, Aberdeen, Elean Gamhna, Alps, Yugoslavia, Germany, Norway, Pyrenees I, Pyrenees II, North Spain, and South Spain.

Details

Corbet et al. (1970) report a study of water voles (genus *Arvicola*) in which the aim was to compare British populations of these animals with those in Europe, to investigate whether more than one species might be present in Britain. The original data consisted of observations of the presence or absence of 13 characteristics in about 300 water vole skulls arising from six British populations and eight populations from the rest of Europe. The data are the percentage incidence of the 13 characteristics in each of the 14 samples of water vole skulls.

Source

G. B. Corbet, J. Cummins, S. R. Hedges, W. J. Krzanowski (1970), The taxonomic structure of British water voles, genus *Arvicola*. *Journal of Zoology*, **61**, 301–316.

Examples

```
data("watervoles", package = "HSAUR")
watervoles
```

waves

Electricity from Wave Power at Sea

Description

Measurements of root mean square bending moment by two different mooring methods.

Usage

```
data("waves")
```

Format

A data frame with 18 observations on the following 2 variables.

method1 Root mean square bending moment in Newton metres, mooring method 1

method2 Root mean square bending moment in Newton metres, mooring method 2

Details

In a design study for a device to generate electricity from wave power at sea, experiments were carried out on scale models in a wave tank to establish how the choice of mooring method for the system affected the bending stress produced in part of the device. The wave tank could simulate a wide range of sea states and the model system was subjected to the same sample of sea states with each of two mooring methods, one of which was considerably cheaper than the other. The question of interest is whether bending stress differs for the two mooring methods.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("waves", package = "HSAUR")
plot(method1 ~ method2, data = waves)
```

weightgain

Gain in Weight of Rats

Description

The data arise from an experiment to study the gain in weight of rats fed on four different diets, distinguished by amount of protein (low and high) and by source of protein (beef and cereal).

Usage

```
data("weightgain")
```

Format

A data frame with 40 observations on the following 3 variables.

source source of protein given, a factor with levels Beef and Cereal.

type amount of protein given, a factor with levels High and Low.

weightgain weight gain in grams.

Details

Ten rats are randomized to each of the four treatments. The question of interest is how diet affects weight gain.

Source

D. J. Hand, F. Daly, A. D. Lunn, K. J. McConway and E. Ostrowski (1994). *A Handbook of Small Datasets*, Chapman and Hall/CRC, London.

Examples

```
data("weightgain", package = "HSAUR")
interaction.plot(weightgain$type, weightgain$source,
                weightgain$weightgain)
```

womensrole

Womens Role in Society

Description

Data from a survey from 1974 / 1975 asking both female and male responders about their opinion on the statement: Women should take care of running their homes and leave running the country up to men.

Usage

```
data("womensrole")
```

Format

A data frame with 42 observations on the following 4 variables.

education years of education.

sex a factor with levels Male and Female.

agree number of subjects in agreement with the statement.

disagree number of subjects in disagreement with the statement.

Details

The data are from Haberman (1973) and also given in Collett (2003). The questions here are whether the response of men and women differ.

Source

S. J. Haberman (1973), The analysis of residuals in cross-classified tables. *Biometrics*, **29**, 205–220.

D. Collett (2003), *Modelling Binary Data*. Chapman and Hall / CRC, London. 2nd edition.

Examples

```
data("womensrole", package = "HSAUR")
summary(subset(womensrole, sex == "Female"))
summary(subset(womensrole, sex == "Male"))
```

Index

* datasets

agefat, 2
aspirin, 3
BCG, 4
birthdeathrates, 5
bladdercancer, 6
BtheB, 7
clouds, 8
CYGOB1, 9
epilepsy, 10
Forbes2000, 11
foster, 12
gardenflowers, 13
GHQ, 13
heptathlon, 14
Lanza, 17
mastectomy, 18
meteo, 19
orallesions, 20
phosphate, 20
pistonrings, 21
planets, 22
plasma, 23
polyps, 24
polyps3, 25
pottery, 26
rearrests, 27
respiratory, 27
roomwidth, 28
schizophrenia, 29
schizophrenia2, 30
 schooldays, 31
skulls, 32
smoking, 33
students, 34
suicides, 35
toothpaste, 35
voting, 36
water, 37

watervoles, 38
waves, 39
weightgain, 40
womensrole, 41

* misc

HSAURtable, 16

agefat, 2
aspirin, 3

BCG, 4
birthdeathrates, 5
bladdercancer, 6
BtheB, 7

clouds, 8
CYGOB1, 9

dist, 13

epilepsy, 10

Forbes2000, 11
foster, 12

gardenflowers, 13
GHQ, 13

heptathlon, 14
HSAURtable, 16

Lanza, 17

mastectomy, 18
meteo, 19

orallesions, 20

phosphate, 20
pistonrings, 21
planets, 22
plasma, 23

polyps, [24](#), [25](#)

polyps3, [25](#)

pottery, [26](#)

rearrests, [27](#)

respiratory, [27](#)

roomwidth, [28](#)

schizophrenia, [29](#)

schizophrenia2, [30](#)

schooldays, [31](#)

skulls, [32](#)

smoking, [33](#)

students, [34](#)

suicides, [35](#)

table, [20](#), [21](#), [27](#), [35](#)

toLatex, [16](#), [17](#)

toLatex.dftab (HSAURtable), [16](#)

toLatex.tabtab (HSAURtable), [16](#)

toothpaste, [35](#)

voting, [36](#)

water, [37](#)

watervoles, [38](#)

waves, [39](#)

weightgain, [40](#)

womensrole, [41](#)